

Pebbling on $C_{4k+3} \times G$ and $M(C_{2n}) \times G$ *

Zheng-Jiang Xia, Yong-Liang Pan[†], Jun-Ming Xu

School of Mathematical Sciences,
University of Science and Technology of China,
Hefei, Anhui, 230026, P. R. China
Email: ylp@ustc.edu.cn

Abstract: The pebbling number of a graph G , $f(G)$, is the least p such that, however p pebbles are placed on the vertices of G , we can move a pebble to any vertex by a sequence of moves, each move taking two pebbles off one vertex and placing one on an adjacent vertex. It is conjectured that for all graphs G and H , $f(G \times H) \leq f(G)f(H)$. If the graph G satisfies the odd two-pebbling property, we will prove that $f(C_{4k+3} \times G) \leq f(C_{4k+3})f(G)$ and $f(M(C_{2n}) \times G) \leq f(M(C_{2n}))f(G)$, where C_{4k+3} is the odd cycle of order $4k+3$ and $M(C_{2n})$ is the middle graph of the even cycle C_{2n} .

Keywords: Graham's conjecture, cycle, middle graph, pebbling number.

1 Introduction

Pebbling in graphs was first introduced by Chung [2]. Consider a connected graph with a fixed number of pebbles distributed on its vertices. A pebbling move consists of the removal of two pebbles from a vertex and the placement of one pebble on an adjacent vertex. The pebbling number of a vertex v , the target vertex, in a graph G is the smallest number $f(G, v)$ with the property that, from every placement of $f(G, v)$ pebbles on G , it is possible to move one pebble to v by a sequence of pebbling moves. The *t-pebbling number* of v in G is defined as the smallest number $f_t(G, v)$ such that from every placement of $f_t(G, v)$ pebbles, it is possible to move t pebbles to v . Then the *pebbling number* and the *t-pebbling number* of G are the smallest numbers, $f(G)$ and $f_t(G)$, such that from any placement of $f(G)$ pebbles or $f_t(G)$ pebbles, respectively, it is possible to move one or t pebbles, respectively, to any specified target vertex by a sequence of pebbling moves. Thus, $f(G)$ and $f_t(G)$ are the maximum values of $f(G, v)$ and $f_t(G, v)$ over all vertices v .

*Supported by "the Fundamental Research Funds for the Central Universities" and the NSF of the People's Republic of China(Grant No. 61272008, No. 11271348 and No. 10871189).

[†]Corresponding author: ylp@ustc.edu.cn

Chung [2] defined the two-pebbling property of a graph, and Wang [9] extended her definition to the odd two-pebbling property as follows.

Suppose p pebbles are located on G , let k be the number of occupied vertices (vertices with at least one pebble), and r be the number of vertices with an odd number of pebbles. G satisfies the *two-pebbling property* means two pebbles can be moved to any vertex of G whenever $p > 2f(G) - k$, and the *odd two-pebbling property* means two pebbles can be moved to any vertex of G whenever $p > 2f(G) - r$. It is clear that the graph which satisfies the two-pebbling property also satisfies the odd two-pebbling property, and there exist graphs do not satisfy the two-pebbling property, which are called the Lemke graphs.

The *middle graph* of a graph G , denoted by $M(G)$, is obtained from G by inserting a new vertex into each edge of G , and joining the new vertices by an edge if the two edges they inserted share the same vertex of G . For any two graphs G and H , we define the *Cartesian product* $G \times H$ to be the graph with vertex set $V(G \times H)$ and edge set the union of $\{((a, v), (b, v)) | (a, b) \in E(G), v \in E(H)\}$ and $\{((u, x), (u, y)) | u \in V(G), (x, y) \in E(H)\}$.

The following conjecture [2], by Ronald Graham, suggests a constraint on the pebbling number of the product of two graphs.

Conjecture 1.1 (Graham) *The pebbling number of $G \times H$ satisfies $f(G \times H) \leq f(G)f(H)$.*

There are a few results verify Graham's conjecture. It has been proved that Graham's conjecture holds for a tree by a graph with the odd-two-pebbling property, especially a tree by a tree [6], an even cycle by a graph with the odd two-pebbling property, a cycle by a cycle [5], a complete graph by a graph with the two-pebbling property [2] and a complete bipartite graph by a graph with the two-pebbling property [3], a fan graph by a fan graph and a wheel graph by a wheel graph [4], a thorn graph of the complete graph with every $p_i > 1$ by a graph with the two-pebbling property [10], and the middle graph of an odd cycle by the middle graph of a cycle [11].

In Section 2, we show that Graham's conjecture holds for the odd cycle C_{4k+3} by a graph with the odd two-pebbling property.

In Section 3, we show that Graham's conjecture holds for the middle graph of an even cycle by a graph with the odd two-pebbling property, especially, the middle graph of an even cycle by the middle graph of an even cycle.

Given a distribution of pebbles on G , Let $p(K)$ be the number of pebbles on a subgraph K of G , $p(v)$ be the number of pebbles on vertex v of G and $k(K)$ ($r(K)$) to be the number of vertices of K with at least one pebble (with an odd number of pebbles). Moreover, denote by $\tilde{p}(K)$ and $\tilde{p}(v)$ the number of pebbles on K and v after some sequence of pebbling moves, respectively.

Let T be a tree with a specified vertex v . T can be viewed as a directed tree denoted by $\vec{T}_v$ with edges directed toward a specified vertex v , also called the root. A

path-partition is a set of non-overlapping directed paths the union of which is $\vec{T}_v$. The path-size sequence of a path-partition $P_1, \dots, P_n$, is an n -tuple $(a_1, \dots, a_n)$, where a_j is the length of P_i (i.e., the number of edges in it), with $a_1 \geq a_2 \geq \dots \geq a_n$. A path-partition is said to majorize another if the nonincreasing sequence of its path size majorizes that of the other. That is, $(a_1, a_2, \dots, a_r) > (b_1, b_2, \dots, b_t)$ if and only if $a_i > b_i$ where $i = \min\{j : a_j \neq b_j\}$. A path-partition of a tree T is said to be maximum if it majorizes all other path-partitions.

The following two Lemmas will be the key tools in the next sections.

Lemma 1.2 ([2]) *The pebbling number $f_t(T, v)$ for a vertex v in a tree T is $t2^{a_1} + 2^{a_2} + \dots + 2^{a_r} - r + 1$, where $a_1, a_2, \dots, a_r$ is the sequence of the path sizes in a maximum path-partition of $\vec{T}_v$.*

Lemma 1.3 ([6]) *If T is a tree, and G satisfies the odd two-pebbling property, then $f(T \times G, (x, g)) \leq f(T, x)f(G)$ for every vertex g in G . In particular, if $P_m = x_1x_2 \dots x_m$ be a path, then $f(P_m \times G, (x_i, g)) \leq f(P_m, x_i)f(G) = (2^{i-1} + 2^{m-i} - 1)f(G) \leq 2^{m-1}f(G)$.*

2 $C_{4k+3} \times G$

In 2003, D. S. Herscovici proved the following two theorems about cycles.

Theorem 2.1 ([5]) *If G satisfies the odd two-pebbling property, then*

$$f(C_{2n} \times G) \leq f(C_{2n})f(G) = 2^n f(G).$$

Theorem 2.2 ([5]) *Suppose G is a graph with $m \geq 5$ vertices which satisfies the odd two-pebbling property and the following inequality*

$$4f_4(G) < 14f(G) - 2(m - 5), \quad (2.1)$$

then, $f(C_{2n+1} \times G) \leq f(C_{2n+1})f(G)$ for $n \geq 3$.

The inequality (2.1) holds for all odd cycles, but does not hold even for paths or even cycles. In this section, we show the following theorem.

Theorem 2.3 *If G satisfies the odd two-pebbling property, then*

$$f(C_{4k+3} \times G) \leq f(C_{4k+3})f(G).$$

Throughout this section, we use the following notation.

Let the vertices of C_{4k+3} be $\{v_0, v_1, \dots, v_{4k+1}, v_{4k+2}\}$ in order. We define the vertex subsets A and B of C_{4k+3} by

$$A = \{v_1, v_2, \dots, v_{2k}\}, B = \{v_{2k+3}, v_{2k+4}, \dots, v_{4k+2}\}$$

For simplicity, among $C_{4k+3} \times G$, let $p_i = p(v_i \times G)$, $r_i = r(v_i \times G)$, $p(A) = p(A \times G)$, $p(B) = p(B \times G)$. Thus, the number of pebbles in a distribution on $C_{4k+3} \times G$ is given by $p_0 + p(A) + p(B) + p_{2k+1} + p_{2k+2}$.

Lemma 2.4 ([7]) *The pebbling numbers of the odd cycles C_{4k+1} and C_{4k+3} are*

$$f(C_{4k+1}) = \frac{2^{2k+2} - 1}{3} = 1 + 2^2 + 2^4 + \dots + 2^{2k}.$$

$$f(C_{4k+3}) = \frac{2^{2k+3} + 1}{3} = 1 + 2^1 + 2^3 + \dots + 2^{2k+1}.$$

Lemma 2.5 *Let $P_{2k} = x_1x_2 \dots x_{2k}$ be a path with length $2k - 1$, and let g be some vertex in a graph G which satisfies the odd two-pebbling property. Then, from any arrangement of $(2^1 + 2^3 + \dots + 2^{2k-1})f(G)$ pebbles on $P_{2k} \times G$, it is possible to put a pebble on every (x_i, g) at once, where $i = 1, 3, \dots, 2k - 1$.*

Proof. We use induction on k , where the case $k = 1$ is trivial.

Suppose that there are $(2^1 + 2^3 + \dots + 2^{2k-1})f(G)$ pebbles on $P_{2k} \times G$. Then there are at least $(2^1 + 2^3 + \dots + 2^{2k-3})f(G)$ pebbles on $\{x_3, x_4, \dots, x_{2k}\} \times G$ (or on $\{x_1, x_2, \dots, x_{2k-2}\} \times G$). By induction, we can use these pebbles to put one pebble to each of these vertices $\{(x_3, g), (x_5, g), \dots, (x_{2k-1}, g)\}$ (or $\{(x_1, g), (x_3, g), \dots, (x_{2k-3}, g)\}$). By Lemma 1.2, $f(P_{2k}, x_1) = 2^{2k-1}$, $f(P_{2k}, x_{2k-1}) = 2^{2k-2} + 1 \leq 2^{2k-1}$. Thus with the remaining $2^{2k-1}f(G)$ pebbles, one pebble can be moved to (x_1, g) (or (x_{2k-1}, g)), and we are done. $\blacksquare$

Similarly, we can get the following Lemma.

Lemma 2.6 *Let $P_{2k+1} = x_1x_2 \dots x_{2k+1}$ be a path with length $2k$, and let g be some vertex in a graph G which satisfies the odd two-pebbling property. Then, from any arrangement of $(2^2 + 2^4 + \dots + 2^{2k})f(G)$ pebbles on $P_{2k+1} \times G$, it is possible to put a pebble on every (x_i, g) at once, where $i = 1, 3, \dots, 2k - 1$.*

From the proof of Theorem 3.2 in [5], it follows that

Lemma 2.7 ([5]) *If $p(A) \geq 2^{2k-1}f(G)$, then with $f(C_{4k+3})f(G)$ pebbles on $C_{4k+3} \times G$, one pebble can be moved to (v_0, g) .*

Proof of Theorem 2.3:

Suppose that there are $f(C_{4k+3})f(G)$ pebbles located on $C_{4k+3} \times G$, then

$$p_0 + p_{2k+1} + p_{2k+2} + p(A) + p(B) = (1 + 2^1 + 2^3 + \dots + 2^{2k+1})f(G). \quad (2.2)$$

Without loss of generality, we may assume that $p(A) \geq p(B)$ and the target vertex is (v_0, g) . The case $k = 0$ is trivial, so we assume that $k \geq 1$.

Note that the vertices of $B \cup \{v_{2k+2}\} \cup \{v_0\}$ form a path isomorphic to P_{2k+2} . It follows from Lemma 1.3 that if we move as many as possible pebbles from $v_{2k+1} \times G$ to $v_{2k+2} \times G$, then one pebble could be moved to (v_0, g) unless

$$\frac{p_{2k+1} - r_{2k+1}}{2} + p_{2k+2} + p(B) + p_0 < 2^{2k+1}f(G). \quad (2.3)$$

From Lemma 2.7, we could move one pebble to (v_0, g) unless

$$p(A) < 2^{2k-1}f(G). \quad (2.4)$$

If (2.3) holds, then

$$\frac{p_{2k+1} + r_{2k+1}}{2} + p(A) > (1 + 2^1 + 2^3 + \dots + 2^{2k-1})f(G). \quad (2.5)$$

From (2.4) and (2.5), we can get $p_{2k+1} + r_{2k+1} > 2f(G)$, and

$$\frac{p_{2k+1} - (2f(G) - r_{2k+1} + 2)}{2} + p(A) \geq (2^1 + 2^3 + \dots + 2^{2k-1})f(G).$$

This implies that we can move enough pebbles from $v_{2k+1} \times G$ to $A \times G$ so that the number of the pebbles on $A \times G$ will reach to $(2^1 + 2^3 + \dots + 2^{2k-1})f(G)$, and at the same time there are h_{2k+1} pebbles are kept on $v_{2k+1} \times G$, where

$$h_{2k+1} = \begin{cases} 2f(G) - r_{2k+1} + 2, & \text{if } r_{2k+1} \geq 2, \\ 2f(G), & \text{if } r_{2k+1} \leq 1. \end{cases}$$

Assume that $2x$ pebbles are taken away from $v_{2k+1} \times G$ such that there are x pebbles reaching $A \times G$, i.e.,

$$x + p(A) = (2^1 + 2^3 + \dots + 2^{2k-1})f(G). \quad (2.6)$$

Step 1. With the h_{2k+1} pebbles on $v_{2k+1} \times G$, we can move one pebble to (v_{2k}, g) .

Now there are $p_{2k+1} - 2x - h_{2k+1}$ pebbles on $v_{2k+1} \times G$, namely,

$$\begin{aligned} \tilde{p}_{2k+1} &= p_{2k+1} - 2x - h_{2k+1} \\ &= p_{2k+1} + 2p(A) - (2^2 + \dots + 2^{2k})f(G) - h_{2k+1}. \end{aligned}$$

So the remaining pebbles on $\{v_0, v_{2k+1}, v_{2k+2}\} \times G$ is

$$\begin{aligned} & p_0 + p_{2k+2} + \tilde{p}_{2k+1} \\ &= p_0 + p_{2k+2} + p_{2k+1} + 2p(A) - (2^2 + \dots + 2^{2k})f(G) - h_{2k+1} \\ &\geq p_0 + p_{2k+2} + p_{2k+1} + p(A) + p(B) - (2^2 + \dots + 2^{2k})f(G) - h_{2k+1} \\ &\geq (1 + 2^2 + 2^4 + \dots + 2^{2k})f(G). \end{aligned}$$

For $p_0 < f(G)$ (otherwise one pebble can be moved to (v_0, g) , and we are done), so

$$p_{2k+2} + \tilde{p}_{2k+1} \geq (2^2 + 2^4 + \dots + 2^{2k})f(G). \quad (2.7)$$

Step 2. It follows from (2.6) and Lemma 2.5 that with $(2^1 + 2^3 + \dots + 2^{2k-1})f(G)$ pebbles on $A \times G$, we can put one pebble to each vertex of $\{(v_1, g), (v_3, g), \dots, (v_{2k-1}, g)\}$.

Step 3. From the inequality (2.7) and Lemma 2.6, it follows that, with $(2^2 + 2^4 + \dots + 2^{2k})f(G)$ pebbles on $\{v_{2k+1}, v_{2k+2}\} \times G$, we can put one pebble to each vertex of $\{(v_2, g), (v_4, g), \dots, (v_{2k}, g)\}$.

The above three steps implies that at least one pebble can be moved to (v_0, g) . ■

3 $M(C_{2n}) \times G$

Throughout this section, we will use the following notations (see Fig.1).

Let $C_{2n} = v_0v_1 \cdots v_{2n-1}v_0$. The middle graph of C_{2n} , denoted by $M(C_{2n})$, is obtained from C_{2n} by inserting u_i into $v_iv_{(i+1) \bmod (2n)}$, and connecting $u_iu_{(i+1) \bmod (2n)}$ ($0 \leq i \leq 2n-1$). The graph $M^*(C_{2n})$ is obtained from $M(C_{2n})$ by removing the edges v_iu_i for $1 \leq i \leq n-1$, $u_{n-1}u_n$ and u_jv_{j+1} for $n \leq j \leq 2n-2$.

We define the vertex subsets A and B of $V(M^*(C_{2n}))$ by

$$A = \{v_1, v_2, \dots, v_{n-1}, u_0, u_1, \dots, u_{n-1}\},$$

$$B = \{v_{n+1}, v_{n+2}, \dots, v_{2n-1}, u_n, u_{n+1}, \dots, u_{2n-1}\}.$$

For simplicity, among $M(C_{2n}) \times G$ (or $M^*(C_{2n}) \times G$), let $p_i = p(v_i \times G)$, $r_i = r(v_i \times G)$, $q_i = p(u_i \times G)$, $s_i = r(u_i \times G)$, $p(A) = p(A \times G)$, $p(B) = p(B \times G)$.




Figure 1: The graphs $M(C_6)$ and $M^*(C_6)$.

Lemma 3.1 ([6]) *Trees satisfy the two-pebbling property.*

Lemma 3.2 ([11]) *If $n \geq 2$, then $f(M(C_{2n})) = 2^{n+1} + 2n - 2$.*

From Lemma 1.2 and the proof of Lemma 3.2, it is not hard to obtain the following

Lemma 3.3 *If $n \geq 2$, then $f(M^*(C_{2n}), v_0) = 2^{n+1} + 2n - 2$.*

Proposition 3.4 *$M(C_{2n})$ satisfies the two-pebbling property.*

Proof. Since symmetry, it is clear that

$$f(M(C_{2n})) = \max\{f(M(C_{2n}), v_0), f(M(C_{2n}), u_0)\}.$$

Assume that the target vertex is v_0 , and $p + k \geq 2f(M(C_{2n})) + 1$. Since $k \leq 4n \leq f(M(C_{2n}))$, we have $p \geq f(M(C_{2n})) + 1$. Thus if there is one pebble located on v_0 , then

with the remaining $f(M(C_{2n}))$ pebbles, a second pebble can be moved to v_0 .

Now, suppose that $p(v_0) = 0$. We will prove that with the same arrangement of pebbles on $M^*(C_{2n})$, two pebbles can be moved to v_0 .

Let $H = M^*(C_{2n})$, $C = H[A \setminus v_1]$, and $D = H[B \setminus v_{2n-1}]$. Then by Lemma 1.2,

$$f(C) = f(D) = 2^{n-1} + n - 2,$$

$$f(C \cup \{v_0\}) = f(D \cup \{v_0\}) = 2^n + n - 2,$$

$$f(C \cup \{v_n\}) = f(D \cup \{v_n\}) = 2^n + n - 2.$$

We consider the worst case, that is $p(v_1) = k(v_1) = p(v_{2n-1}) = k(v_{2n-1}) = 1$ (where $k(v_i) = 1$ if there is at least one pebble located on v_i and 0 otherwise), then

$$p(C) + k(C) + p(D) + k(D) + p_n + q_n + 4 \geq 2^{n+2} + 4n - 3.$$

If $p(C) + k(C) > 2^{n+1} + 2n - 4$, then by Lemma 3.1, two pebbles can be moved to v_0 . Thus we may assume that $p(C) + k(C) \leq 2^{n+1} + 2n - 4$ and $p(D) + k(D) \leq 2^{n+1} + 2n - 4$. We will show that every one of the vertices u_0 and u_{2n-1} will get at least two pebbles by a sequece of pebbling moves.

Let $p'_n = 2^{n+1} + 2n - 4 - p(C) - k(C) \geq 0$, and paint all the pebbles on C red along with the p'_n pebbles on v_n . Similarly, paint the pebbles on D black, along with $p''_n = 2^{n+1} + 2n - 4 - p(D) - k(D)$ pebbles on v_n . It is easy to see there are enough pebbles on v_n to do this.

Now either $p(C) + k(C) = 2^{n+1} + 2n - 4$ or there are red pebbles on v_n . If equality holds, then $p(C) \geq 2^n + n - 2$, then two red pebbles can be moved to u_0 . If there are red pebbles on v_n , then $k'_n = 1$, and the red pebbles satisfy

$$p(C) + k(C) + p'_n + k'_n = 2^{n+1} + 2n - 3,$$

and again two red pebbles can be moved to u_0 . Similarly, two black pebbles can be moved to u_{2n-1} , so we can move one red pebble and one black pebble to v_0 .

If the target vertex is u_0 , then a similar argument can show that there are at least two pebbles can be moved to u_0 . ■

Lemma 3.5 ([5]) *Let $P_k = x_1x_2 \dots x_k$ be a path, and let g be some vertex in a graph G which satisfies the odd two-pebbling property. Then, from any arrangement of $(2^k - 1)f(G)$ pebbles on $P_k \times G$, it is possible to put a pebble on every (x_i, g) at once ($1 \leq i \leq k$).*

Lemma 3.6 *Let $P_k = x_1x_2 \dots x_k$ be a path ($k \geq 2$), and g be some vertex in a graph G which satisfies the odd two-pebbling property. Then from any arrangement of $(2^k - 2)f(G)$ pebbles on $x_k \times G$, it is possible to put a pebble on every (x_i, g) at once ($1 \leq i \leq k - 1$).*

Proof. We use induction on k , where the case $k = 2$ is trivial. If it is true for $k - 1$, suppose there are $(2^k - 2)f(G)$ pebbles on $x_k \times G$, we use $(2^{k-1} - 2)f(G)$ pebbles to put a pebble on every (x_i, g) at once ($2 \leq i \leq k - 1$), and with the remaining $2^{k-1}f(G)$ pebbles we can put one pebble on (x_1, g) . ■

Lemma 3.7 *Let T_k be the graph obtained from P_k by joining x_i to a new vertex y_i ($1 \leq i \leq k-1$), where $P_k = x_1x_2 \dots x_k$ is a path ($k \geq 2$). Let g be some vertex in a graph G which satisfies the odd two-pebbling property. Then for any arrangement of $(2^k + k - 3)f(G)$ pebbles on $T_k \times G$, one of the following will occur*

- (1) *we can put a pebble on every (x_i, g) at once ($1 \leq i \leq k-1$);*
- (2) *we can put two pebbles on (x_1, g) .*




Figure 2: The graph T_k in Lemma 3.7.

Proof. While $k = 2$, by Lemma 1.2 and Lemma 1.3, with $3f(G)$ pebbles on $T_2 \times G$, one pebble can be moved to the vertex (x_1, g) .

Suppose that there are $(2^k + k - 3)f(G)$ pebbles on $T_k \times G$ for $k \geq 3$. Let $T'_k = T_k \setminus \{x_1, y_1\}$. Clearly, $T'_k \cong T_{k-1}$.

If $p(T'_k \times G) < (2^{k-1} + k - 4)f(G)$, then $p(P_{x_1y_1} \times G) \geq (2^{k-1} + 1)f(G) \geq 5f(G)$. Then clearly, we can move two pebbles to (x_1, g) .

If $p(T'_k \times G) \geq (2^{k-1} + k - 4)f(G)$, and $p_k \geq (2^{k-1} - 2)f(G)$. By Lemma 3.6, using $(2^{k-1} - 2)f(G)$ pebbles on $x_k \times G$, we can put a pebble on every (x_i, g) for $2 \leq i \leq k-1$. With the remaining $(2^{k-1} + k - 1)f(G)$ pebbles, by Lemma 1.2 and Lemma 1.3, we can put one pebble on (x_1, g) for $f(T_k \times G, (x_1, g)) \leq f(T_k, x_1)f(G) = (2^{k-1} + k - 1)f(G)$.

If $p(T'_k \times G) \geq (2^{k-1} + k - 4)f(G)$, and $p_k < (2^{k-1} - 2)f(G)$. We use induction in this case, while the case $k = 2$ holds.

Let r_y be the number of vertices with an odd number of pebbles in $\{y_2, y_3, \dots, y_{k-1}\} \times G$. We only need to take off r_y pebbles from $\{y_2, y_3, \dots, y_{k-1}\} \times G$ so that each vertex in it has an even number of pebbles. It is clear that $r_y \leq (k-2)|V(G)| \leq (k-2)f(G)$, so $r_y + p_k < (2^{k-1} + k - 4)f(G)$. So we can choose $(2^{k-1} + k - 4)f(G)$ pebbles from $T'_k \times G$ which contains all pebbles on $x_k \times G$, so that the number of the remaining pebbles on each vertex of $\{y_2, y_3, \dots, y_{k-1}\} \times G$ is even except at most one vertex. By induction, with these $(2^{k-1} + k - 4)f(G)$ pebbles we can put one pebble on every (x_i, g) at once for $2 \leq i \leq k-1$ or move two pebbles to (x_2, g) and then at least one pebble can be moved to (x_1, g) .

Now we prove that with the remaining $(2^{k-1} + 1)f(G)$ pebbles, one pebble can be moved to (x_1, g) .

Let $\tilde{p}_y = \sum_{i=2}^{k-1} \tilde{p}(y_i \times G)$. Let P_1 denote the path $y_1x_1x_2 \dots x_{k-1}$, and P_2 denote the path $x_1x_2 \dots x_{k-1}$.

For the number of the remaining pebbles on each vertex of $\{y_2, y_3, \dots, y_{k-1}\} \times G$ is even except at most one vertex, then we can move $\lfloor \frac{1}{2}\tilde{p}_y \rfloor$ pebbles from the vertices of $\{y_2, y_3, \dots, y_{k-1}\} \times G$ to $\{x_2, x_3, \dots, x_{k-1}\} \times G$.

Case 1. $\tilde{p}_y \leq 2^{k-1}f(G) - 1$. Then

$$\tilde{p}(P_1 \times G) = (2^{k-1} + 1)f(G) - \tilde{p}_y + \left\lfloor \frac{1}{2}\tilde{p}_y \right\rfloor \geq (2^{k-2} + 1)f(G).$$

By Lemma 1.3, $f(P_1 \times G, (x_1, g)) \leq f(P_1, x_1)f(G) = (2^{k-2} + 1)f(G)$, so one pebble can be moved to (x_1, g) .

Case 2. $\tilde{p}_y \geq 2^{k-1}f(G)$. Then $\tilde{p}(P_2 \times G) \geq \lfloor \frac{1}{2}\tilde{p}_y \rfloor \geq 2^{k-2}f(G)$. By Lemma 1.3, $f(P_2 \times G, (x_1, g)) \leq f(P_2, x_1)f(G) = 2^{k-2}f(G)$, so one pebble can be moved to (x_1, g) . ■

Theorem 3.8 *If G satisfies the odd two-pebbling property, then*

$$f(M(C_{2n}) \times G) \leq f(M(C_{2n}))f(G) = (2^{n+1} + 2n - 2)f(G).$$

Proof. Suppose that there are $(2^{n+1} + 2n - 2)f(G)$ pebbles placed on the vertices of $M(C_{2n}) \times G$, we will show that at least one pebble can be moved to the target vertex.

Since symmetry, it is clear that

$$f(M(C_{2n}) \times G) = \max\{f(M(C_{2n}) \times G, (v_0, g)), f(M(C_{2n}) \times G, (u_0, g))\}.$$

So we only need to distinguish two cases.

Case 1. The target vertex is (v_0, g) .

Subcase 1.1. $p_n + r_n \leq 2f(G)$.

We remove all the pebbles off $v_n \times G$ such that

$$\begin{aligned} \tilde{p}((M^*(C_{2n}) \setminus v_n) \times G) &= \frac{p_n - r_n}{2} + p(A) + p(B) + p_0 \\ &= -\frac{1}{2}(p_n + r_n) + p_n + p(A) + p(B) + p_0 \\ &\geq (2^{n+1} + 2n - 3)f(G). \end{aligned}$$

By Lemma 1.2, $f(M^*(C_{2n}) \setminus v_n, v_0) = 2^{n+1} + 2n - 3$. According to Lemma 1.3, one pebble can be moved to (v_0, g) .

Subcase 1.2. $p_n + r_n > 2f(G)$.

Then we can put two pebbles to (v_n, g) . Note that p_n and r_n are of the same-parity, we keep $2f(G) - r_n + 2$ pebbles on $v_n \times G$ so that at least two pebbles still can be moved to (v_n, g) , and move the rest pebbles to $A \times G$. So

$$\tilde{p}(A \times G) = \frac{1}{2}(p_n - (2f(G) - r_n + 2)) + p(A) = \frac{p_n + r_n}{2} - f(G) - 1 + p(A). \quad (3.1)$$

By Lemma 1.2, $f(M^*(C_{2n})[B, v_0], v_0) = 2^n + n - 1$, so if we move as many as possible pebbles from $v_n \times G$ to $B \times G$, then one pebble can be moved to (v_0, g) unless

$$\frac{p_n - r_n}{2} + p(B) + p_0 \leq (2^n + n - 1)f(G) - 1. \quad (3.2)$$

If (3.2) holds, then

$$\begin{aligned} \tilde{p}(A \times G) &= \frac{1}{2}(p_n + r_n) - f(G) - 1 + p(A) \\ &\geq p_n + p(A) + p(B) + p_0 - f(G) - (2^n + n - 1)f(G) \\ &= (2^{n+1} + 2n - 2)f(G) - (2^n + n)f(G) \\ &= (2^n + n - 2)f(G). \end{aligned} \quad (3.3)$$

It follows from Lemma 1.2 that

$$f(M^*(C_{2n}) \setminus \{v_n, u_{n-1}, v_{n-1}\}, v_0) = 3 \cdot 2^{n-1} + 2n - 4.$$

Thus if we move as many as possible pebbles from $v_n \times G$ to $u_n \times G$, and from $v_{n-1} \times G$ to $u_{n-2} \times G$, then one pebble can be moved to (v_0, g) unless

$$\begin{aligned} \frac{1}{2}(p_n - r_n) + \frac{1}{2}(p_{n-1} - r_{n-1}) + p(B) + p_0 + (p(A) - p_{n-1} - q_{n-1}) \\ \leq (3 \cdot 2^{n-1} + 2n - 4)f(G) - 1. \end{aligned} \quad (3.4)$$

If (3.4) holds, then $\frac{1}{2}(p_n + r_n) + \frac{1}{2}(p_{n-1} + r_{n-1}) + q_{n-1} \geq (2^{n-1} + 2)f(G) + 1$.
Thus

$$\left(\frac{1}{2}(p_n + r_n) - f(G) - 1 + q_{n-1}\right) + \left(\frac{1}{2}(p_{n-1} + r_{n-1}) - f(G) - 1\right) \geq 2^{n-1}f(G) - 1. \quad (3.5)$$

Subcase 1.2.1. $\frac{1}{2}(p_n + r_n) - f(G) - 1 + q_{n-1} \geq f(G)$.

Then from (3.3) it follows that, with $f(G)$ pebbles on $u_{n-1} \times G$, one pebble can be moved to (u_{n-1}, g) ; and from Lemma 3.7 it follows that, with the remaining $(2^n + n - 3)f(G)$ pebbles, we can put one pebble to each (u_i, g) for $0 \leq i \leq n - 2$ or put two pebbles to (u_0, g) , we can move one more pebble to (u_{n-1}, g) with $2f(G) - r_n + 2$ pebbles on $v_n \times G$, so one pebble can be moved to (v_0, g) .

Subcase 1.2.2. $\frac{1}{2}(p_n + r_n) - f(G) - 1 + q_{n-1} < f(G)$.

Then from (3.5), we have

$$\frac{p_{n-1} + r_{n-1}}{2} - f(G) - 1 \geq (2^{n-1} - 1)f(G). \quad (3.6)$$

So we can keep $2f(G) - r_{n-1} + 2$ pebbles on $v_{n-1} \times G$, so that one pebble can be moved to (u_{n-2}, g) , and moving no less than $(2^{n-1} - 1)f(G)$ pebbles to $u_{n-2} \times G$. With these pebbles, by Lemma 3.5, we can put one pebble to every (u_i, g) at once ($0 \leq i \leq n - 2$). So one pebble can be moved to (v_0, g) .

Case 2. The target vertex is (u_0, g) .

Let $M'(C_{2n})$ be the graph obtained from $M(C_{2n})$ by removing the edges $u_i v_{i+1}$ for $0 \leq i \leq n - 2$ and $u_j v_j$ for $n + 2 \leq j \leq 2n - 1$ and $u_n v_n, u_n v_{n+1}, u_0 v_0$.

Let $A' = \{u_1, u_2, \dots, u_{n-1}, v_1, v_2, \dots, v_n\}$ and $B' = \{u_{n+1}, u_{n+2}, \dots, u_{2n-1}, v_{n+1}, v_{n+2}, \dots, v_{2n-1}, v_0\}$.

It is clear that $M'(C_{2n})[A'] \cong M^*(C_{2n})[A]$, and $M'(C_{2n})[B', u_0] \cong M^*(C_{2n})[B, v_0]$. We only need to prove that one pebble can be moved from $M'(C_{2n}) \times G$ to (u_0, g) .

Subcase 2.1. $q_n + s_n \leq 2f(G)$. By a similar process as before, one pebble can be moved to (u_0, g) .

Subcase 2.2. $q_n + s_n > 2f(G)$.

Then by a similar process as before, we can keep $2f(G) - s_n + 2$ pebbles on $u_n \times G$ so that two pebbles can be moved to (u_n, g) , and move the rest pebbles to $A' \times G$. So

$$\begin{aligned} \tilde{p}(A' \times G) &= \frac{q_n - (2f(G) - s_n + 2)}{2} + p(A') \\ &= \frac{q_n + s_n}{2} - f(G) - 1 + p(A'). \end{aligned}$$

Similarly, if we move as many as possible pebbles from $u_n \times G$ to $B' \times G$, then one pebble can be moved from $B' \times G$ to (u_0, g) , unless

$$\tilde{p}(A' \times G) \geq (2^n + n - 2)f(G).$$

According to Lemma 3.7, we can put one pebble on (u_i, g) at once for $1 \leq i \leq n - 1$ or put two pebbles on (u_1, g) . With $2f(G) - s_n + 2$ pebbles on $u_n \times G$, one more pebble can be moved to (u_{n-1}, g) . So one pebble can be moved to (u_0, g) . ■

4 Remark

In this paper, we show that $f(C_{4k+3} \times G) \leq f(C_{4k+3})f(G)$, where G satisfies the odd two-pebbling property. But we cannot show $f(C_{4k+1} \times G) \leq f(C_{4k+1})f(G)$. Maybe since the odd cycles have not been solved completely, the middle graph of the odd cycle is an open problem, too.

References

- [1] J. Akiyama, T. Hamada, I. Yoshimura, Miscellaneous properties of middle graphs, *TRU Math*, **10** (1974), 41-53.
- [2] F. R. K. Chung, Pebbling in hypercubes, *SIAM J. Discrete Math.*, **2**(4) (1989), 467-472.
- [3] R. Feng, J. Y. Kim, Graham's pebbling conjecture of production complete bipartite graph, *Sci. China Ser. A*, **44** (2001), 817-822.
- [4] R. Feng, J. Y. Kim, Pebbling numbers of some graphs, *Sci. China Ser. A*, **45** (2002), 470-478.
- [5] D. Hersovici, Grahams conjecture on products of cycles, *J. Graph Theory*, **42** (2003), 141-154.
- [6] D. Moews, Pebbling Graphs, *J. Combin. Theory, Ser. B*, **55** (1992), 244-252.
- [7] L. Pachter, H.S. Snevily, B. Voxman, On pebbling graphs, *Congr. Numer.*, 107(1995), 65-80.
- [8] H. S. Snevily, J. A. Foster, The 2-pebbling property and conjecture of Graham's, *Graphs Combin.*, **16** (2000), 231-244.
- [9] S. S. Wang, Pebbling and Graham's conjecture, *SIAM J. Discrete Math.*, **226**(1-3) (2001), 431-438.
- [10] Z. P. Wang, Y. Zou, H. Liu, Z. Wang, Graham's pebbling conjecture on product of thorn graphs of complete graphs, *SIAM J. Discrete Math.*, **309**(10) (2009), 3431-3435.
- [11] Y. S. Ye, F. Liu, M. Q. Zhai, Pebbling numbers of middle graphs of cycles and Graham's conjecture, *Operations Research Transactions*, **17**(3) (2013), 35-44.